



III NICA DAYS 2019

International scientific and engineering conference associated with the
IVth MPD Collaboration Meeting
and
V Slow Control Warsaw 2019

21- 25 October 2019



Ministry of Science
and Higher Education
Republic of Poland

Warsaw University
of Technology

Correlation femtoscopy at the NICA energies

P. Batyuk, L. Malinina, K. Mikhaylov, G.Nigmatkulov


on behalf of PWG

October 22, 2019

Outline:

- **Femtoscopy and Motivation**
- **Hybrid vHLLE+UrQMD model**
- **Comparison with STAR BES**
- **First look at factorial moments with vHLLE+UrQMD**
- **???**
- **Other activities we are responsible for**

Femtoscscopy formalism



Correlation femtoscopy:

Measurement of space-time characteristics R , c_τ of particle production using particle correlations due to the effects of quantum statistics (QS) and final state interactions (FSI)

Two-particle correlation function:

theory: $C(q) = \frac{N_2(p_1, p_2)}{N_1(p_1)N_2(p_2)}$, $C(\infty) = 1$

experiment: $C(q) = \frac{S(q)}{B(q)}$, $q = p_1 - p_2$

$S(q)$ is a distribution of pair momentum difference of particles from the same event

$B(q)$ is a reference distribution built by mixing of particles from different events

Parametrizations used:

1D CF:

$$C(q_{inv}) = 1 + \lambda e^{-R^2 q_{inv}^2}$$

R is a Gaussian radius in PRF,

λ is a correlation strength parameter

1D-analysis is sensitive only to the system size averaged over all directions.


3D CF:

$$C(q_{out}, q_{side}, q_{long}) = 1 + \lambda e^{-R_{out}^2 q_{out}^2 - R_{side}^2 q_{side}^2 - R_{long}^2 q_{long}^2}$$

Both R and q are in Longitudinally Co-Moving Frame (LCMS)

3D-analysis gives an access to the three system sizes in three directions separately.

Definition of femtoscopy radii:




S. Pratt. Phys. Rev. D 33 (1986) 1314

G. Bertsch. Phys. Rev. C37 (1988) 1896

Motivation

- **Femtoscropy allows one:**
 - To obtain spatial and temporal information on particle-emitting source at kinetic freeze-out
 - To study collision dynamics depending on EoS
- **RHIC Beam Energy Scan program (BES-I):** $\sqrt{s_{NN}} = 7.7, 11.5, 19.6, 27, 39$ GeV
Measured pion and kaon femtoscopic parameters:
 m_T -dependences of radii,
flow-induced $x - p$ correlations
- **NICA energy range:** $\sqrt{s_{NN}} = 4 - 11$ GeV

Phys. Rev. C92 (2015) 1, 014904



Femtoscscopy with vHLE+UrQMD

Iu. Karpenko, P. Huovinen, H. Petersen, M. Bleicher,
Phys.Rev. C 91, 064901 (2015)



Parameters τ_0 , R_{\perp} , R_{η} and η/s
adjusted using basic observables

in the RHIC BES-I region.

$\sqrt{s_{NN}}$ [GeV]	τ_0 [fm/c]	R_{\perp} [fm]	R_{η} [fm]	η/s
7.7	3.2	1.4	0.5	0.2
8.8 (SPS)	2.83	1.4	0.5	0.2
11.5	2.1	1.4	0.5	0.2
17.3 (SPS)	1.42	1.4	0.5	0.15
19.6	1.22	1.4	0.5	0.15
27	1.0	1.2	0.5	0.12
39	0.9	1.0	0.7	0.08
62.4	0.7	1.0	0.7	0.08
200	0.4	1.0	1.0	0.08

Model tuned by matching
with existing
experimental data from
SPS and BES-I RHIC

EoS to be used in the
model



- Chiral EoS - crossover transition
J. Steinheimer et al., J. Phys. G 38, 035001 (2011)
- Hadron Gas + Bag Model
1-st order phase transition
P. F. Kolb et al., Phys.Rev. C 62, 054909 (2000)

Hydrodynamic phase lasts
longer with 1PT, especially
at lower energies but
cascade smears this
difference.

Pion emission time

(a) - after hydrodynamic phase


(b) - after cascade




3D Pion radii versus m_T with vHLE+UrQMD

Phys. Rev. C 96, 024911

(2017)




Comparison of extracted radii with the STAR data



Crossover EoS “works” better for lowest collision energies.

- R_{out} (XPT) at high energies and R_{out} (1PT) at all energies are slightly overestimated
- $R_{out, long}$ (1PT) $>$ $R_{out, long}$ (XPT) by value of $\sim 1-2$ fm.


R_{out}/R_{side} with vHLE + UrQMD model



Exp. data:




R_{out}/R_{side} and $R_{out}^2 - R_{side}^2$ as a function of $\sqrt{s_{NN}}$ at a fixed m_T demonstrate a wide maximum near $\sqrt{s_{NN}} \approx 20$ GeV

Our calculations:



R_{out}/R_{side} (XPT) agrees with almost all STAR data points within rather large systematic errors, while R_{out}/R_{side} (1PT) overestimates the data.

Ratio of $R_{out,side,long}(1PT)/R_{out,side,long}(XPT)$ vs. $\sqrt{s_{NN}}$




- R_{side} practically coincide for both scenarios
- R_{out} and R_{long} for 1PT EoS are greater than for XPT EoS demonstrating a strong k_T -dependence


Why?

The difference comes from a **weaker transverse flow developed in the fluid phase** with 1PT EoS as compared to XPT EoS and its **longer lifetime** in 1PT EoS

Pions:




Kaons:



Analysis:

- AuAu, $\sqrt{s_{NN}} = 11.5$ GeV
- $N_{events} \approx 400000$
- Standard 3D Gaussian fit used
- Projections of 3D-kaon correlation functions on out-side-long directions are more Gaussian
- XPT CF projections on long direction are visibly wider than 1PT especially for kaons

Pion and kaon radii vs. m_T with vHLE+UrQMD



Important to measure both kaon and pion radii!

- As well as for pions kaon out and long radii are greater for 1PT than for XPT
- Approximate m_T -scaling for pions and kaons observed only for side radii
- Out almost flat for 1PT
- R_{long} (kaons) is greater than R_{long} (pions) due to larger average time emission
- R_{out} / R_{side} for kaons is less than for pions
- Approximately the same result is for AuAu $\sqrt{s_{NN}} = 7.7$ GeV

Do we put here anything
from our recent studies?
????

Factorial moments with vHLE+UrQMD

Proposed by A. Bialas and R. Peschanski (Nucl. Phys. B 273 (1986) 703) to study the dependence of the normalized factorial moments of the rapidity distribution on the size of the resolution


Set of definitions of moments and cumulants

$$F_i = M^{i-1} \cdot \left\langle \frac{\sum_{j=1}^M k_j \cdot (k_j - 1) \cdot \dots \cdot (k_j - i + 1)}{N \cdot (N - 1) \cdot \dots \cdot (N - i + 1)} \right\rangle$$


- **No variation of moments δy expected** if fluctuations are purely statistical
- **Observation of variations indicates the presence of physics origin fluctuations**

Intermittency (**fluctuations of many different sizes in 1D, 2D and 3D space**) has been studied at LEP, Tevatron, Protvino in ee, hh, hA, AA interactions at various energies.

Factorial moments with vHLE+UrQMD



Factorial moments with $v\text{HLE} + \text{UrQMD}$



Different energy dependence is expected for XPT and 1PT EoS

Other activities we do

vHLE+UrQMD interface software

How to get?

1. `git clone https://github.com/pbatyuk/vHLE_package.git`
2. `git checkout 1.1.2`

How to compile and use?

- `vHLE_package/README.md` (very detailed description on how to ...)

Aim of the project:

- To collect all components (model + interface) in one place.
- To start simulations locally or remotely in a common way.
- To avoid a huge messy in the start configure scripts.
- Possibility to use the model for its adjustment (pre-hydro + hydro phase) as planned.

Main macro: vHLLE_package/macro/vHLLE.C

```

void vHLLE() {
VHLLE* gen = new VHLLE();
gen->SetSourceROOT(""); // Set ROOT-environment if not set yet and necessary to be set
// gen->SetExtendedFileName(kTRUE); // Set use of extended output filename ...
gen->SetUseBatch(kFALSE); // False value (default) means calculations at your locale machine ...
gen->SetBatchCluster("ncx"); // Possible values are: ncx, govorn, basov and gsi

// Parameters below (6) are considered as those to be set obligatory
gen->SetPathToTheModel(""); // Absolute(!) path to the root folder of the model
gen->SetOutputDirectory(""); // Directory where output data stored
gen->SetEnergy(7.7); // Set collision energy [GeV], possible energies are 7.7 GeV ...
gen->SetImpact(0., 3.3); // Set impact range (min, max) [fm]
gen->SetEoS("XPT"); // Set EoS to be used (1PT - first order phase transition, XPT - crossover)
gen->SetNsamples(100); // nEvents to be sampled in hadronic cascade from one hydro-evolution

gen->SetParameters(); // Set parameters for urqmd, hydro and hadronic cascade given by ...

// Modifiers to redefine almost all parameters given by the author for urqmd, hydro ...
// See $VHLLE/vhllc.h to get more if needed
// N. B.: Redefinition, if needed, can be done after gen->SetParameters() called !!!
/*
gen->SetTau0(3.2);
gen->SetEtaS(0.2);
gen->SetRg(1.4);
gen->SetRgz(0.5);
gen->SetNsamples(100);
*/

gen->PrintBasicParams();
gen->CheckParamsValidity(); // It checks whether the params defined are consistent
gen->GenerateStartScript(); // It produces a script to be executed
delete gen;
}

```

Package for Femtoscopic Analysis

Femtoscopy


- Inherited from STAR (StHbtMaker) and ALICE (AliFemto)
- Keeps the same hierarchy as in ALICE (PckgName/, PckgNameUser/, macros/)
- Works with ROOT 5 and 6
- Lighter than ancestors:
 - Most of STAR-developed classes replaced with ROOT ones
 - Better compression, smaller sizes
- Implemented running options (INDEPENDENT on experiment-dependent software):
 - Standalone mode – compile with g++ (clang) and run on your “laptop”
 - Maker; Tasks will be also implemented

Data formats (DST)

- General-purpose data format for Monte Carlo generators - McDst
- Similar to UniGen (developed at GSI)
- Lighter, faster, easy expandable, works with ROOT 5 and 6, g++ (clang)
- Possibility to add converters from other generators: Terminator, EPOS, AMPT ...
- Group has a positive experience on the data format developments:
 - PicoDst format in STAR (standard data format for physics analysis)

Package for Femtoscopic Analysis

Output ROOT tree:



It allows:

- To set track cuts, particle pair cuts, number of events to be used for mixing ...
- To get 1D and 3D correlation functions for a set of k_T -bins
- To switch on / off different physics effects (QS, FSI ...)

Main macro to define conditions of user's analysis

```
int main(int argc, char* argv[]) {
    ...
    // Create and set track cut
    trackCut->setPdgId(particlePdg);
    trackCut->setEta(-1., 1.);
    trackCut->setPt(0.15, 1.55);
    trackCut->setMass(particleMass);
    ...
    // Set how many events to mix
    hbtAnalysis->setNumEventsToMix(10);
    ...
    // Lednicky weight generator
    hbtWeight->setPairType(pairType);
    hbtWeight->setCoulOn();
    hbtWeight->setQuantumOn();
    hbtWeight->setStrongOff();
    hbtWeight->set3BodyOff();
    ...
    // Create 1D correlation function
    // integrated over kT
    StHbtModelQinvCorrFctn *oneDim =
    new StHbtModelQinvCorrFctn
    ("hTheorQinv", 40, 0., 0.4);
    // Create 3D correlation function
    // integrated with kT binning
    StHbtModelBPLCMS3DCorrFctnKt *threeDim =
    new StHbtModelBPLCMS3DCorrFctnKt
    ("hTheorBPLCMS", 80, -0.4, 0.4, 4,
    0.15, 0.59);
}
```

Thank you for attention!